

# Intellectual Fertilizer Prediction System for Precision Agriculture Using Optimized Artificial Networks

Bhagwan Dinkar Thorat<sup>1</sup>, Dr. Sunita A. Jahirabadkar<sup>2</sup>

<sup>1</sup>Research Scholar, Smt. Kashibai Navale COE, SPPU Pune, India

<sup>2</sup>Professor, Cummins College of Engg. Pune, India

<sup>1</sup>bhagwan.thorat@gmail.com

<sup>2</sup>sunita.jahirabadkar@cumminscollege.in

## ARTICLE INFO

## ABSTRACT

Received: 10 Nov 2024

Revised: 28 Dec 2024

Accepted: 14 Jan 2025

In agriculture, effective fertilizer management is essential to increasing crop yields and developing sustainable agricultural practices. Devices with artificial intelligence (AI) have produced agricultural tools that help farmers achieve controlled and accurate farming. To meet farmer expectations, increase yield production, and oversee agricultural operations, it is more crucial to anticipate fertilizer correctly. A unique Automated Fertilizer Management System (AFMS) using an Optimized Artificial Neural Network (OANN) model, augmented by back propagation and chain rule techniques, is presented in this study using deep learning (DL) models. The proposed method refines the back propagation process and increases learning efficiency by utilizing advanced optimization techniques such as regularization and hyper parameter modifying. During training, the model guarantees precise and effective weight updates by utilizing the chain rule for gradient calculation. A wide range of data, including soil nutrient levels and climate parameters unique to the Indian agricultural context, are used to train the ANN. These attributes are then sent into an "Optimized Artificial Neural Network (OANN)" that uses previously acquired data to forecast fertilizer results. Specifically, back propagation and chain rule techniques are used to modify the OANN weights in order to increase the prediction accuracy of the classifier.

**Keywords:** Optimized Artificial Neural Network, backpropagation, Deep Learning, fertilizer, soil nutrient, BPCR

## 1. Introduction

Agriculture is the main source of income for the majority of people in India [16]. A sizable section of the population depends on agriculture for their livelihood, which also boosts the economy. However, the effective use of fertilizers remains a critical challenge for maximizing crop yields and ensuring sustainable farming practices [6]. Conventional fertilizer application techniques frequently result in either excessive or insufficient use, which can have a negative impact on crop output, soil health, and the environment. There is increasing interest in using cutting-edge technologies for accurate fertilizer forecast in order to overcome these issues.

The most crucial element to guaranteeing maximum field productivity is agronomy. Crop production and management comprise a number of processes in the majority of agricultural techniques. In agriculture, turning and loosening the soil is the first step. After the soil has been sufficiently loosened and aerated, it is prepared for crop cultivation. Depending on the soil's requirements, fertilizers and manures are added to raise the soil's nutritional content. However, excessive fertilizer application harms the soil, while insufficient fertilizer causes crop nutrition deficiencies [25]. Using Artificial Neural Networks (ANNs), especially with the backpropagation method, is one potential strategy [17]. This technique makes it possible to model intricate interactions between different crop growth-influencing elements, such as crop types, soil characteristics, and climatic circumstances. The backpropagation technique effectively updates the neural network's weights by utilizing the chain rule, which enhances the neural network's capacity to forecast the ideal fertilizer requirements. Using artificial neural networks

(ANN) to anticipate fertilizer can greatly improve farmers' decision-making in the Indian environment, which has a variety of agroclimatic variables.

Artificial neural network (ANN) models can be taught to make customized fertilizer recommendations by combining data from multiple sources, including soil testing, remote sensing, and historical yield data. In addition to maximizing resource utilization, this strategy encourages sustainable farming methods, which support both environmental preservation and food security. The application of deep learning methods, especially ANN backpropagation, is a viable solution to enhance fertilizer management and improve crop results as India's agricultural environment develops. New prediction techniques utilizing OANN are proposed in order to address the shortcomings of traditional methods and correct the factors influencing the soil. An effective optimization algorithm is also presented. Through the implementation of new, crucial challenges in fertilizer prediction analysis, this study contributes to its certainty. With the least amount of computational complexity, it aids in the establishment of an efficiency model for fertilizer forecast in all meteorological aspects. Deep learning algorithms and emerging and growing features in agronomy are expected to be discovered in this work[15], offering a solid foundation for developing methods in fertilizer prediction algorithms.

## 2. Related Work

Sivasan karan et. al.[1] study examines a suggested model for forecasting the nutrient composition of soil and suggesting the best fertilizers for growing groundnut crops in a sustainable manner. The model is evaluated against current machine learning classifiers and is based on an Enhanced 1D-CNN deep learning methodology. It seeks to enhance generalization, accuracy, and performance while addressing the shortcomings of the existing soil nutrient prediction models. The suggested model performs better than current models when tested utilizing nutrient datasets from a particular region. The significance of soil fertility for crop production and the influence of technology on precision-based agriculture are also highlighted in the document. It also highlights the need of soil nutrients like potassium, phosphorus, and nitrogen for crop growth and talks about the difficulties in predicting nutrients and soil fertility.

Jion Gao et.al.[5] the study, emphasizing the value of swarm intelligence and machine learning in creating efficient fertilization decision models. It provides context for the writers' contributions to the subject by highlighting the necessity of creative methods to maximize fertilizer application for soybean, rice, and maize crops. It also discusses how fertilizer use affects the environment and promotes sustainable methods that reduce harm to ecosystems. In order to support methods that increase output while minimizing environmental damage, the authors situate their study within the larger framework of sustainable agriculture.

Deepak Mane et. al.[7] The study tackles the serious problem of plant infections, which endanger the production of therapeutic herbs, especially basil. To increase the precision of disease identification, the authors suggest a hybrid deep learning model that integrates Convolutional Neural Networks (CNN) with Support Vector Machine (SVM) and K-Nearest Neighbour (KNN) algorithms. A hybrid model for the early identification of diseases in basil plants—which are significant in traditional medicine—is the main goal of this work.

Amilia Nongbet et. al. [18] investigates the potential benefits of applying nano fertilizers in modern farming. Increased crop yield, reduced nutrient losses, and enhanced nutrient uptake efficiency are the goals of nano fertilizers. They offer advantages like regulated nutrient release, targeted nutrition delivery to plants, and less environmental effect.

Takashi S. et.al. [2] explains how to forecast crop yield and make fertilizer recommendations using machine learning models. The study assesses how yield and economically optimal input rates (EOIRs) are affected by various machine learning techniques and covariate selection. It emphasizes the need for caution when using machine learning for fertilizer recommendations by highlighting how sensitive predicted EOIRs and profits are to algorithm and covariate selection. The difficulties with model uncertainty and the significance of taking temporal and regional variability into account when modeling crop yield response are also covered in the study. All things considered, it highlights the necessity of giving careful thought to model uncertainty and the limitations of machine learning techniques in order to produce trustworthy site-specific input management suggestions.

Pratik kr et. al. [9] explains how deep learning models can be used to identify plant leaf deficiency and suggest suitable fertilizers. It describes the data used, the methodology, data pretreatment methods, and the outcomes.

Image processing, classification using the ResNet-50 architecture, and comparing the model's performance to that of other pre-trained models are the main topics of the study. The usefulness of the suggested approach in obtaining high accuracy in identifying malnutrition in photos of plant leaves is highlighted in the text. It also emphasizes how the study could be expanded in the future to anticipate further plant leaf picture classifications.

Dr. T. Venkat Narayana Rao et. al. [17] The use of artificial neural networks (ANN) in agriculture is its main area of interest. The key concepts include employing artificial neural networks (ANN) to estimate agricultural production and soil quality, suggesting appropriate fertilizers, and solving the difficulties faced by farmers. The potential of machine learning to enhance farming methods and lower the suicide incidence among farmers is highlighted in the research. The design of the suggested system and the backpropagation algorithm's application to neural network model training are also covered. Overall, the study emphasizes how important it is to use technology to solve problems in agriculture.

Thilina Abekoon et al. [15] explains the creation of a deep learning model to forecast the concentrations of the three main soil nutrients—potassium, phosphorus, and nitrogen—that are essential for cabbage growth. The algorithm analyses and forecasts soil nutrient content based on plant growth factors. The study describes the process, which includes gathering data and training the model with various transfer functions. The findings show that the deep learning model accurately predicts soil nutrient levels, especially when the tangent sigmoid transfer function is used. With possible uses in sustainable agriculture, the study highlights the importance of this strategy for maximizing fertilizer use and improving soil nutrition.

Vani, et al. [14] created a PLMDC strategy to improve clustering accuracy for both deeply allocated and sparse agricultural large data. To clean the data, a logical linear regression approach is employed. A genetic algorithm (GA) was used to extract features. In order to forecast crop yields based on their characteristics, the A-FP growth algorithm was introduced. This technique improved clustering performance.

### 2.1 Gap Analysis

Fertilizer prediction employs a number of methods and algorithms to forecast fertilizer and overcome obstacles. However, there are still certain disadvantages, such as the need for additional fertilizers, the maximum manpower costs for small-holder farmers, and crop nutrition deficiencies. There were certain difficulties in evaluating different kinds of study. However, The limitations of traditional modeling approaches, the significance of thorough data utilization, the potential of advanced modeling techniques, the need for regionally adaptable models, and the need for more focused research on the relationship between crop yield and fertilization management are all highlighted [5]. The cloud cover must be automated, and image capture is a dangerous procedure. In order to achieve more clustering accuracy with less complexity, the PLMDC scheme was created in [14], and GA was used for the feature extraction genetic algorithm. The crop management approach was offering increased yields along with improved soil aggregation and less soil moisture losses. However, it costs small-holder farmers more in effort and is ineffective for agronomic water management techniques. linear regression (LR) analysis, which boosts soybean yield and lessens the climate impact of crop cultivation; however, it necessitates a lengthy growing season with high thermal requirements.

## 3. Proposed Architecture

In order to predict the kind and amount of fertilizer needed for the best crop growth, the suggested deep learning-based fertilizer prediction model makes use of a multi-input architecture that analyses a variety of agricultural data sources, including soil parameters (such as pH, nitrogen, potassium and phosphorus) and meteorological factors (such as temperature, humidity, and rainfall) [25]. Three essential steps are part of the applied prediction approach: "pre-processing, feature extraction, and fertilizer forecast." The pre-processing step is where the data cleansing procedure is completed. By identifying and removing outliers from the fresh training dataset, the data cleaning method [25] is used to enhance the quality of training data.

These characteristics are then predicted using an optimized ANN, which forecasts the fertilizer as the yield information. The backpropagation chain rule (BPCR) is suggested as a way to adjust the OANN weights during training in order to improve prediction accuracy. The created backpropagation\_chain rule (BPCR) prediction framework is demonstrated in Figure 1.




Fig1. Fertilizer prediction system block diagram

Precision agriculture is made possible by this model, which offers customized fertilizer suggestions that maximize crop output, reduce environmental effect, and conform to local farming methods. It also continuously updates and receives feedback to adjust to changing environmental conditions.

3.1 Dataset Preparation

Dataset is very important factor for using deep learning model in modern agriculture from which dataset we can predict fertilizer accurately. There are total 8 features in dataset out of that 7 features are in-depended and 1 is depended. Total 2230 samples are in dataset. This dataset can be found from various sources such as Agricultural Colleges, Soil Testing Labs, Government websites. We have visited Vasant Dada Sugar Institute Pune, Rajarambapu sugar factory Islampur, Sangliand Agriculture College Karad and collected datasets from them. Also we have generated our own dataset using different sensors.




Fig2. Sample dataset images

N	P	K	pH	EC	Temp	Moisture	Target
138	8.6	560	7.46	0.62	26	38	Urea
213	7.5	338	7.62	0.75	29	45	DAP
163	9.6	718	7.59	0.51	34	62	10-26-26
157	6.8	475	7.64	0.58	32	34	Urea
270	9.9	444	7.63	0.4	28	46	10-26-26
220	8.6	444	7.43	0.65	26	35	Urea
220	7.2	222	7.62	0.43	25	64	DAP
207	7	401	7.63	0.59	33	50	Urea
289	8.6	560	7.58	0.44	30	42	Urea
138	8.1	739	7.55	0.33	29	33	10-26-26

Table 1. Data samples from dataset

### 3.2 Exploratory Data Analysis

Once the data is collected, it needs to be cleaned and transformed into a format that is suitable for feeding into a deep learning model.

#### 3.2.1 Data Cleaning

**Handle Missing Values:** Either remove rows or columns with a high number of missing values, or use imputation (such as substituting the mean or median for missing data). **Finding and handling outliers** that could skew model training is known as outlier detection. Methods such as IQR (Interquartile Range) can be applied. **Noise Reduction:** If sensor data (such as NPK, soil moisture or temperature) is noisy, use smoothing techniques. **Normalization** to scale continuous numerical features such (soil NPK, pH, temperature, and moisture) to fall within a comparable range [0,1] mean ( $\mu_w$ ) of 0 and standard deviation ( $\sigma_w$ ) of 1 use Z-score normalization.

$$Z' = \frac{X - \mu_w}{\sigma_w} \quad (1)$$

In Eq.1  $Z'$  denotes newvalue,  $X$  is the value from the dataset,  $\mu_w$  denotes mean and  $\sigma_w$  denotes standard deviation. One method for transforming category variables into numerical values is label encoding. Label encoding is frequently used for categorical information, such as fertilizer type, in fertilizer prediction. Each distinct value of a categorical feature in label encoding is substituted with an integer, usually beginning with 0 for the first category, 1 for the second, and so forth.

### 4. OANN Model for Fertilizer prediction using Back-Propagation with Chain-Rule algorithm (OANN+BPCR)

An Optimized Artificial Neural Network (OANN) is a kind of neural network model that uses optimization strategies and cutting-edge learning algorithms to produce predictions with high efficiency and accuracy as shown in Fig3. Based on dataset an OANN can be used to predict the type and quantity of fertilizer that will be best appropriate for a particular soil nutrients, and weather circumstances. The Backpropagation Algorithm with the Chain Rule (BPCR) is a powerful technique used in training neural networks, including OANNs, by updating the weights of the network during the training process. This process minimizes the error between the model's predicted output and the actual target output fertilizer requirement using a gradient descent-based optimization technique.




Fig.3 Backpropagation Framework

In the context of agriculture, predicting the type and quantity of fertilizer needed based on various soil and environmental conditions is crucial. It leverages the backpropagation algorithm, employing the chain rule to compute gradients efficiently, thereby updating the network's weights and biases to improve accuracy. By training the model on a dataset of soil and environmental conditions, it learns to predict the appropriate fertilizer type or quantity needed.

Let's define the input features for the fertilizer prediction problem as follows:

$X = [x_1, x_2, \dots, x_n]$  is input vector representing soil properties, weather conditions, etc.  $x_1$ -Nitrogen(N),  $x_2$ -Phosphorus(P),  $x_3$ -Potassium(K),  $x_4$ -Soil pH,  $x_5$ -Temperature,  $x_6$ -Humidity. To predict the fertilizer denoted as  $y_{\text{pred}}$ , which could be the fertilizer type. Consider a Artificial neural network with the Input layer Consists of  $n$  features  $x_1, x_2, \dots, x_n$ . Hidden layer(s) contains  $h$  neurons, output layer produces a prediction  $y_{\text{pred}}$ . Let the weights between the input layer and the hidden layer be denoted by  $W_1$  and the biases by  $b_1$ . For the hidden layer, the output can be expressed as Eq.2

$$z_1^{(i)} = \sum_{j=1}^n W_1^{(i,j)} x_j + b_1^{(i)} \quad (2)$$

Where  $i$  represents the  $i^{\text{th}}$  neuron in the hidden layer.  $W_1^{(i,j)}$  are the weights connecting the  $j^{\text{th}}$  input feature to the  $i^{\text{th}}$  hidden neuron.  $b_1^{(i)}$  is the bias term for the  $i^{\text{th}}$  hidden neuron.  $z_1^{(i)}$  is the weighted sum of inputs to the  $i^{\text{th}}$  hidden neuron. The activation function transforms the weighted sum in Eq.3

$$a_1^{(i)} = f(z_1^{(i)}) \quad (3)$$

where  $f(z)$  is a non-linear sigmoid activation function ( $f(z) = \frac{1}{1+e^{-z}}$ ). Similarly, for the output layer with weights  $W_2$  and bias  $b_2$  Eq.4

$$z_2 = \sum_{i=1}^h W_2^{(i)} a_1^{(i)} + b_2 \quad (4)$$

The final output  $y_{\text{pred}}$  is Eq.5

$$y_{\text{pred}} = f_{\text{output}}(z_2) \quad (5)$$

where  $f_{\text{output}}$  is the activation function for the output layer

The neural network's ability to estimate the amount of fertilizer needed is measured by the OANN's loss function. OANN loss function was indicated as loss and is denoted the Eq. 6

$$L = \frac{1}{N} \sum_{k=1}^N (y_{\text{pred},k} - y_{\text{true},k})^2 \quad (6)$$

where  $N$  is the number of training samples.  $y_{\text{pred},k}$  is the predicted fertilizer value for sample  $k$ .  $y_{\text{true},k}$  is the true fertilizer requirement for sample  $k$ . Here, the OANN weights with the best performance are selected through backpropagation algorithm using the chain rule method. To reduce the loss  $L$ , the weights  $W_1$  and  $W_2$  are updated by backpropagation. The gradients of the loss function with respect to each weight are calculated throughout the update phase using the chain rule.

### Step 1: Determine the Output Layer Gradients:

In relation to the output layer, the gradient of the loss  $L$  is Eq.7

$$\frac{\partial L}{\partial z_2} = \frac{\partial L}{\partial y_{\text{pred}}} \cdot \frac{\partial y_{\text{pred}}}{\partial z_2} \quad (7)$$

so  $y_{\text{pred}} = f_{\text{output}}(z_2)$  we have Eq. 8

$$\frac{\partial y_{\text{pred}}}{\partial z_2} = f'_{\text{output}}(z_2) \quad (8)$$

Consequently, the output layer's gradient is represent the Eq. 9

$$\frac{\partial L}{\partial z_2} = (y_{\text{pred}} - y_{\text{true}}) \cdot f'_{\text{output}}(z_2) \quad (9)$$

Next, Eq. 10 is determine the gradients in relation to the output layer's weights

$$\frac{\partial L}{\partial W_2^{(i)}} = \frac{\partial L}{\partial z_2} \cdot a_1^{(i)} \quad (10)$$

Eq. 11 is calculate the output layer's bias gradient

$$\frac{\partial L}{\partial b_2} = \frac{\partial L}{\partial z_2} \quad (11)$$

### Step 2: Determine the Hidden Layer's Gradients:

Now, Eq. 12 we backpropagate to the hidden layer using the chain rule, the gradient of the loss with respect to the activations of the hidden layer neurons

$$\frac{\partial L}{\partial a_1^{(i)}} = \frac{\partial L}{\partial z_2} \cdot W_2^{(i)} \cdot f'_1(z_1^{(i)}) \quad (12)$$

Where  $f'_1(z_1^{(i)})$  is the derivative of the activation function used in the hidden layer? Then, compute the gradient with respect to the weights in the hidden layer Eq. 13

$$\frac{\partial L}{\partial W_1^{(ij)}} = \frac{\partial L}{\partial a_1^{(i)}} \cdot x_j \quad (13)$$

And the Eq. 14 is bias gradient for the hidden layer:

$$\frac{\partial L}{\partial b_1^{(i)}} = \frac{\partial L}{\partial a_1^{(i)}} \quad (14)$$

### Step 3: Update the Weights:

Finally, the weights are updated using an optimization algorithm, such as **Gradient Descent**:

$$W_1^{(i,j)} \leftarrow W_1^{(i,j)} - \eta \cdot \frac{\partial L}{\partial W_1^{(i,j)}} \quad (15)$$

$$W_2^{(i)} \leftarrow W_2^{(i)} - \eta \cdot \frac{\partial L}{\partial W_2^{(i)}} \quad (16)$$

$$b_1^{(i)} \leftarrow b_1^{(i)} - \eta \cdot \frac{\partial L}{\partial b_1^{(i)}} \quad (17)$$

$$b_2 \leftarrow b_2 - \eta \cdot \frac{\partial L}{\partial b_2} \quad (18)$$

where  $\eta$  is the learning rate. After training the model through multiple iterations (epochs), the optimized weights  $W_1$  and  $W_2$  are used to predict the fertilizer requirements for new input data:

$$y_{\text{pred}} = f_{\text{output}} \left( \sum_{i=1}^h W_2^{(i)} a_1^{(i)} + b_2 \right) \quad (19)$$

where  $a_1^{(i)}$  is the output of the hidden layer neurons, computed using the learned weights and activation functions.

This mathematical model describes how an artificial neural network (ANN) uses backpropagation with the chain rule to forecast fertilizer needs. After specifying the input features, the weighted sums at each layer are calculated, activation functions are applied, the loss is calculated, and the weights are updated using the chain rule to minimize the loss. Through iterative weight optimization,

## 5. Investigational Results

Python was used to implement a recommended fertilizer prediction method utilizing the OANN+BPCR model. The fertilizer prediction in this work, as previously indicated. This covers the soil's temperature, moisture content, nutrients, and a few other characteristics. Furthermore, the suggested model's efficacy was assessed in comparison to conventional models with learning rates of 70, 80, and 90. Furthermore, it was established that the OANN+BPCR model that was presented in terms of error analysis of MAE, MSE, RMSE, and RMAE. Performance was assessed using a dataset of prepared soil nutrients. In order to predict fertilizer, various models (OANN+BPCR, Ensemble, RF, KNN, and SVM) were applied to the soil nutrients dataset. Fig. 5 the OANN+BPCR based sigmoid function had the highest accuracy of 95.02% when compared to all other models. Fig.4 shows the training and validation loss performance graph. The validation accuracy is quite low at first, but it tends to grow as the number of epochs increases. After a given number of epochs, the value tends to reach saturation.




Fig.4 Validation and Training Performance graph

The overall error analysis of the approved OANN+BPCR model compared to traditional schemes for the dataset. Error analysis was therefore performed for measures such as MAE, MSE, RMSE, and RMAE. The accepted OANN+BPCR model has achieved minimal error (MAE: 0.306837) when compared to existing methods shown in Fig.6.




Fig.5 Overall Accuracy Graph




Fig.6 Error Analysis Graph

## 6. Conclusion

This study used the OANN+BPCR algorithm to create a new fertilizer prediction model. Prior to deriving the "soil parameters, metrological parameters, statistical features, and higher order statistical features," preprocessing was completed. The characteristics were then provided as input to OANN for prediction, where the OANN+BPCR model was used to fine-tune the OANN weights. The provided system was ultimately found to be superior to the conventional schemes in a number of aspects. For all learning rates, the error output of the proposed method mainly reaches minimal values. The most widely used acceptable scheme has a learning rate of 70 and a minimum MAE value of 0.247. Stated otherwise, the created OANN+BPCR technique outperformed 95.02% the existing SVM, RF, KNN, and Ensemble (SVM+RF+KNN) models by 72%, 78.4%, 82.5%, and 85.4% when learning rates were 70. Consequently, the efficacy of the introduced OANN+BPCR technique effectively established its superiority. Some of the constraints of the proposed research include its inability to manage large amounts of fertilizer data, also known as big data or high dimensional data. Furthermore, a few restrictions, including soil and metrological factors, can be used to anticipate fertilizer. Therefore, the study will be expanded in the future to include various meteorological conditions or soil factors related to micronutrients. To increase the system's accuracy, research can concentrate on investigating different deep learning models or implementing more sophisticated strategies.

## References

- [1] Sivasan karan S, Dr. K. Jagan Mohan , Dr. G. Mohammed Nazer" Soil Nutrients Prediction and Optimal Fertilizer Recommendation for Sustainable Cultivation of Groundnut Crop using Enhanced-1DCNN DLM " International Journal of Advanced Computer Science and Applications (IJACSA), Vol. 13, No. 4, 2022.
- [2] Takashi S. T. Tanaka, Gerard B. M. Heuvelink, Taro Mieno, David S. Bullock " Can machine learning models provide accurate fertilizer recommendations? " Precision Agriculture Springer 2024.
- [3] Marwan Albahar "A Survey on Deep Learning and Its Impact on Agriculture: Challenges and Opportunities" MDPI 2023
- [4] Hirushan Sajindra, Thilina A bekoon, J.A.D.C.A. Jayakody, UpakaRathnayake," A novel deep learning model to predict the soil nutrient levels (N, P, and K) in cabbage cultivation." Elsevier 2023.
- [5] Jian Gao, Wenzhi Zeng, Zhipeng Ren, Chang Ao, Guoqing Lei, Thomas Gaiser and Amit Kumar Srivastava. "A Fertilization Decision Model for Maize, Rice, and Soybean Based on Machine Learning and Swarm Intelligent Search Algorithms" Agronomy MDPI 2023.
- [6] Bhagwan D. Thorat and Dr. Sunita. A. Jahirabadkar, " Smart Farming in the Indian Context: Fertilizer Prediction through Ensemble Machine Learning Based Soil Nutrient Analysis." Panamerican Mathematical Journal 2023.
- [7] Deepak Mane, Mahendra Deore, Rashmi Ashtagi, Sandip Shinde, Yogesh Gurav " Basil plant leaf disease detection using amalgam based deep learning models " Journal of Autonomous Intelligence (2024).
- [8] S. Bhanumathi, M. Vineeth and N. Rohit "Crop Yield Prediction and Efficient use of Fertilizers" IEEE 2019.
- [9] Prathik Kr, Madhuri Javaregowda, Bhanushree K J "9 Deep Learning-Based Fertilizer Recommendation System Using Nutrient Deficiency Prediction " CSITSS IEEE 2023.
- [10] Jairos Rurinda , Shamie Zingorea , Jibrin M. Jibrin , Tesfaye Balemi, Kenneth Masuki , Jens A. Andersson, Mirasol F. Pampolino , Ibrahim Mohammed , James Mutegi. "Science-based decision support for formulating crop fertilizer recommendations in sub-Saharan Africa." Alpha Agricultural systems 180 (2020), Elsevier.
- [11] Sirsat M. S.,Cernadas,E., Fernandez-Delgado M., & Khan R., "Classification of agricultural soil parameters in India". In Computers and Electronics in Agriculture, Elsevier, vol.135, 269-279 2017.
- [12] Amit Kumar Srivastava, Nima Safaei, Saeed Khaki, Gina Lopez, Wenzhi Zeng, Frank Ewert, Thomas Gaiser & Jaber Rahimi, "Winter wheat yield prediction using convolutional neural networks from environmental and phenological data", scientific reports, 12:3215 ,2022
- [13] Seungtaek Jeong, Jonghan Ko2, Taehwan Shin, Jong-min Yeom, "Incorporation of machine learning and deep neural network approaches into a remote sensing-integrated crop model for the simulation of rice growth" , Scientific Reports ,2022.
- [14] Vani P Suvitha, Rathi S "Improved data clustering methods and integrated A-FP algorithm for crop yield prediction" .Distributed and Parallel Databases:2021.
- [15] Hirushan Sajindra, Thilina Abekoon, J.A.D.C.A. Jayakody, Upaka Rathnayake "A novel deep learning model to predict the soil nutrient levels (N, P, and K) in cabbage cultivation" Smart Agricultural Technology Elsevier 2023.
- [16] Bhagwan D. Thorat and Sunita A. Jahirabdkar "Survey on Precision Agriculture in Indian Context for Effective Fertigation Using Learning Techniques" ICICT 2022 Springer.
- [17] Dr. T. Venkat Narayana Rao, S. Manas "Artificial Neural Networks for Soil Quality and Crop Yield Prediction using Machine Learning" International Journal on Future Revolution in Computer Science & Communication Engineering 2019.
- [18] X.E. Pantazi , D. Moshou , T. Alexandridis , R.L. Whetton , A.M. Mouazen , "Wheat yield prediction using machine learning and advanced sensing Techniques" Computers and Electronics in Agriculture 121 (2016) 57-65
- [19] D. Elavarasan, P.D. Vincent, "Crop yield prediction using deep reinforcement learning model for sustainable agrarian applications", IEEE Access, 8 , pp. 86886-86901, 2020.
- [20] Priti Jorvekar, Sharmila Wagh, Jayshree Prasad," A Comprehensive Review on Deep learning Techniques for Crop Yield Prediction"IEEE 2023..
- [21] M.S. Suchithra \*, Maya L. Pai, "Improving the prediction accuracy of soil nutrient classification by optimizing extreme learning machine parameters" Elsevier 2019..

- 
- [22] Amilia Nongbet, Awdhesh Kumar Mishra, Yugal Kishore Mohanta , SaurovMahanta ,Manjit Kumar Ray , Maryam Khan , Kwang-Hyun Baek and Ishani Chakrabartty, "Nanofertilizers: A Smart and Sustainable Attribute to Modern Agriculture" MDPI 2022.
  - [23] M.T. Fathi, M. Ezziyyani, M. Ezziyyani, S. El Mamoune, "Crop yield prediction using deep learning in Mediterranean Region", International Conference on Advanced Intelligent Systems for Sustainable Development, Springer, 2019
  - [24] Bhuvaneswari Swaminathan, Saravanan Palani, Subramaniaswamy Vairavasundaram "Feature Fusion Based Deep Neural Collaborative Filtering Model For Fertilizer Prediction." Elsevier 2023.
  - [25] Priti Prakash Jorvekar · Sharmila Kishor Wagh · Jayashree Rajesh Prasad "Crop yield predictive modeling using optimized deep convolutional neural network: An automated crop management system". Springer 2023.
  - [26] Sonal Agarwal and SandhyaTatar "A Hybrid Approach For Crop Yield Prediction Using Machine Learning And Deep Learning Algorithms" CONSILIO 2020.
  - [27] P. Sumathi, V. V. Karthikeyan, M. S. Kavitha and S. Karthik "Improved Soil Quality Prediction Model Using Deep Learning for Smart Agriculture Systems " Tech Science Press CSSE 2023.
  - [28] Hammouch, H.; El-Yacoubi, M.; Qin, H.; Berrahou, A.; Berbia, H.; Chikhaoui, M. A two-stage deep convolutional generative adversarial network-based data augmentation scheme for agriculture image regression tasks. In Proceedings of the 2021 International Conference on Cyber-Physical Social Intelligence (ICCSI), Beijing, China, 18–20 December 2021.
  - [29] Sourav, A.I.; Emanuel, A.W.R. Recent Trends of Big Data in Precision Agriculture: A Review. In IOP Conference Series: Materials Science and Engineering; IOP Publishing: Bristol, UK, 2021; Volume 1096, p. 012081.
  - [30] Liu, J.; Cheng, L.; Yan, S. Deep learning with noisy labels. In Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition, Long Beach, CA, USA, 16–20 June 2019.